

Studies on Littorinidae from the Atlantic

BY

KLAUS BANDEL

Institut für Paläontologie der Rheinischen Friedrichs-Wilhelms-Universität, D 53 Bonn, West Germany

(5 Plates; 22 Text figures)

INTRODUCTION

MEMBERS OF THE FAMILY Littorinidae occur as large populations on many shores ranging from the southern Caribbean Sea to the North Atlantic Ocean. Most littorinid species are restricted to hard rock substrates within or above the normal intertidal zone. Few, like *Littorina angulifera* Lamarck, 1822 and *L. nebulosa* (Lamarck, 1822), prefer wood substrates, and only *L. littorea* Linnaeus, 1758 can be found on all substrates, even on muddy bottoms. All 18 species described herein live primarily on plant material which is scraped and bitten from the substrate by the radula. ROSEWATER (1970) in light-microscope studies of many radulae indicated that the more generalized taenioglossate radula of this family offers little promise for systematic diagnosis below the generic level. Here it is demonstrated that with the scanning electron microscope clear differences in the radular morphology of 18 species of Littorinidae can be shown and used to distinguish between species. Furthermore, possible taxonomic relations can be shown more clearly and problems concerning species differentiation in the *L. ziczac* (Gmelin, 1791) and *Nodilittorina tuberculata* (Menke, 1828) groups can be solved.

In defining subgenera in the family Littorinidae the shape of egg capsules should also be considered as a valuable aid, as shown here and as previously suggested by BORKOWSKY & BORKOWSKY (1969), MARCUS & MARCUS (1963), and ROSEWATER (1970).

All individuals used in this study were collected by myself to avoid confusion about the ecological data. Material was collected from Cape Cod (Massachusetts, U. S. A.), 1965; Nassau in the Bahama Islands, 1970; Santa Marta, Colombia from 1970 to 1972; Curaçao (Netherlands Antilles) in 1971; Fuerteventura on the Canary Islands in 1972, Banyuls sur Mer (southern France), 1973; and the Wadden-Sea of the Oosterschelde in the Netherlands.

More detailed taxonomic studies especially connected with the Caribbean species are omitted in this work and will be reported separately (BANDEL & KADOLSKY, in preparation).

DESCRIPTION OF THE ENVIRONMENT

I. Santa Marta (Figures 1 and 2)

Detailed collection and observation over a period of 18 months was carried out between the fall of 1970 and the spring of 1972 in the vicinity of Santa Marta, Colombia.



Figure 1

Transect of a cliff in the Bay of Santa Marta (Isla Morito)

The species that lives under driest conditions of all littorinids and of all marine prosobranchs in this area is *Tectarius muricatus* (Linnaeus, 1758). Individuals of this species are found in greatest abundance on flat benches of rocky shores and cliffs in areas just barely reached by spray of high seas and from strong gales. *Tectarius muricatus* has to endure long periods (months) of dryness. Members of this species can be found only rarely in the bay of Santa Marta on the islands Moro and Morito (Figure 1). Here they occupy the highest platforms carved out of the metamorphic shales by salt erosion. Only in the period between September and May, when strong winds blow almost continuously, can these animals be moistened with sea water spray and can feed on minute algae that grow on the wet rock surface. During the remainder of the year spray only rarely will reach these upper levels, and therefore all individuals remain motionless.

A thriving population of *Tectarius muricatus* is found in the Ensenada Playa Brava north of Santa Marta. Here the platform of a bench consisting of indurated calcareous dune sand receives considerable spray from comparatively heavy wave action. But here also the animals have to endure high temperatures and extreme dryness during long periods of quiet water. Some animals may even be surrounded by crusts of salt remaining from former spray water puddles, but they renew activity after being made wet by fresh sea water.

An unusual occurrence of a *Tectarius muricatus* population was observed in the mangrove belt of the southern extension of the Ensenada Chengue, north of Playa Brava. In this unusually quiet part of the bay the animals live on driftwood and dead branches piled up and anchored between the roots of living mangrove bushes and on a dry shingle beach a few tenths of a centimeter above normal water level. But here also rare high waves caused by extreme swells and unusual wind directions disturb the bay water so that waves go high enough to wet the habitat. Only then they can feed.

On rocky shores below the *Tectarius muricatus* populations *Littorina* sp. and *Nodilittorina tuberculata* populations live with a narrow to wide zone of intermixture. While *N. tuberculata* prefers flat bench surfaces with spray water basins and puddles as substrate, *Littorina* sp. is also likely to be found on vertical rock faces. The mollusks do not cling to the dry rock with their feet, but the lip of the shell is attached to it by a hard and brittle film of mucus. The strength of this film is sufficient to hold the snail to a vertical surface over long periods of time. *Nodilittorina tuberculata* will, like *T. muricatus*, survive encrustation in salt formed by the residue from evaporated spray water. Both species (*L. sp.* and *N. tuberculata*) will withstand extreme dryness and heat, for some places

where they live are moistened only during certain periods of the year. The darker *Littorina* sp. avoids some of the heat of the sun by hiding in crevices and solution cavities which are especially abundant in the supratidal cliff sections. The populations of both species extend from areas of very dry conditions down into areas which are more frequently washed over by sea water. Periodically, the latter environment will not be reached by the tide and dry out. The animals are then exposed to the heat of the tropical sun. Larger pools situated in this area will periodically attain high temperatures (over 40° C) or dry out completely. Changes in the salinity of the water are large and range from saline brines (due to evaporation) to almost fresh water (due to rain). Smaller individuals of *N. tuberculata* and *Littorina* sp. prefer pools closer to the high tide line having a normal or a slightly higher salinity. When rain dilutes this pool water considerably, the juvenile animals leave it and congregate at the rim of the pool above the water surface, but they remain on a wet substrate.

Littorina ziczac (Gmelin, 1791) populations will usually be found in areas where splash and spray are common all year around and below the adult populations of *Littorina* sp. and *Nodilittorina tuberculata*. This habitat coincides with that described for this species from Margarita Island, Venezuela by RODRIGUEZ (1959). On cliffs and pebble beaches with little wave action *L. ziczac* and *L. sp.* form mixed populations, while *N. tuberculata* usually is absent. Here *L. ziczac* is also found more commonly in crevices on the shady side of larger pebbles, but *L. sp.* is common in places exposed to the sun. It can be concluded that *L. ziczac*, in general, needs more moisture even than *L. sp.* although it can endure weeks of dryness exposed to the sun. On rock cliffs more exposed to wave action where spray may moisten up to 10 m of vertical rock surface above the high tide mark, the zone of *L. ziczac* populations is distinct from that of *L. sp.* populations. Here on the island of Morito, for example (Figure 1), a 1 m to ½ m zone of mixed populations is followed by 2 m to 3 m of pure populations above and below, respectively.

The lower range of *Littorina ziczac* populations in the whole area of Santa Marta is always distinct and coincides with the upper limit of the occurrence of *Purpura patula* (Linnaeus, 1758) in the upper splash zone. Often this muricid was seen feeding on littorinids. In places where *P. patula* populations were removed, the *L. ziczac* population expanded down to the high water line in a short time.

In rocky environments, the representative of the Littorinidae which is found in the lowest situation is the very small *Littorina meleagris* Potiez & Michaud, 1838. Indi-

viduals of this species may be found in moderately warm splash pools together with the juveniles of the 3 species *L. ziczac*, *L. sp.*, and *Nodilittorina tuberculata* if water is exchanged often by splash or high waves. The animal cannot tolerate dryness. From shallow pools or depressions which are frequently renewed with sea water to a few centimeters below the water line, *L. meleagris* is very abundant, occurring in greatest numbers in the narrow normal intertidal area (only about 20 cm in height in the vicinity of Santa Marta). Common predators for *L. meleagris* are representatives of the genera *Thais* and *Leucozonia*. Pebble beaches with little wave action are densely settled by *L. meleagris* which are on and below rocks in the intertidal area and among rocks densely covered by algal growths just below low tide mark.




Figure 2

Transect through the estuary of the Cienaga Grande Lagoon between Santa Marta and Barranquilla (Colombia)

Littorina angulifera populations were found near *Tectarius muricatus* settlements only at the unusual locality of the Chengue mangrove belt. Here *L. angulifera* settles on living roots and branches of the mangrove bushes, while *T. muricatus* uses dead branches and washed up driftwood between bushes for attachment.

Littorina angulifera was collected on mangrove bushes growing under normal marine conditions and in brackish water lagoons. Fresh water conditions were tolerated only

for short periods. This was observed in the mouth of the Cienaga Grande, a large, shallow, brackish water lagoon in the Rio Magdalena delta about 40 km south of Santa Marta. Here, fresh river water from a Rio Magdalena flood had in the winter of 1970/1971 flushed out the brackish water of the lagoon. After about 2 months of fresh water flood all the rich settlements of *L. angulifera* had disappeared from the mangrove bushes of the lagoon and its estuary, even though the animals populate the bushes in heights up to 2 m above the water surface (even at flood times with raised water level). Shorter periods of low salinity should be tolerated by this species. In the same Rio Magdalena flood the animals in another lagoon were exposed for at least one month to fresh water; most of these individuals survived.


The zonation of *Littorina angulifera* populations in the vicinity of Santa Marta is the same as described by COOMANS (1969) for the Caribbean region in general, by MARCUS & MARCUS (1963) for Brasil, and by LENDERKING (1954) for Florida.

At the mouth of the Cienaga Grande the environments of *Littorina angulifera* and *L. nebulosa* approach one another. *Littorina angulifera* settles on wooden pilings of an old bridge and *L. nebulosa* can be found next to it on rocks of a breakwater, covered by algae (Figure 2). This is not an unusual environment for *L. nebulosa*, but it demonstrates its tolerance to salinity changes in this estuarine milieu of the lagoon entrance. During the extended periods of freshwater outflow of the Rio Magdalena flood *L. nebulosa* populations disappeared from the estuarine area and survived only on rock piles at the seaward end of the entrance to it. The more usual habitat of *L. nebulosa* is driftwood tree trunks firmly fixed between boulders or anchored on sandy beaches within the surf zone. Strong wave action is tolerated and dense populations are usually found. All littorinids mentioned previously gather microorganisms, algae and detritus from the surface on which they live. *Littorina nebulosa* seems to be able to feed on rotting wood. Therefore, cellulose wood fibres are the principal constituent of their faeces. This is in contrast to *L. angulifera* and *Tectarius muricatus* which may also be collected from driftwood or wooden pilings, but never

Explanation of Figures 23 to 29

Figure 23: *Littorina littorea* from Cape Cod, Massachusetts. × 240
 Figure 24: *Tectarius muricatus* from Paradise Island, Nassau. × 480
 Figure 25: *Tectarius muricatus* from Santa Marta. × 480
 Figure 26: *Littorina saxatilis* from the Oosterschelde. × 350

Figure 27: *Littorina saxatilis* from the Oosterschelde. × 310
 Figure 28: *Littorina obtusata* from the Oosterschelde. × 245
 Figure 29: *Littorina obtusata* from Cape Cod, Massachusetts. × 360



produce faeces consisting primarily of wood fibers (BANDEL, in press).

II. Curaçao (Figure 3)

Near the old wrak in Cornelisbaai on the leeward side of the island of Curaçao, Netherland Antilles, the beach region consists of a hard beach rock bench and limestone boulders. The seaward edge of the rock bench drops suddenly into deeper water where coral growth is evident. The transect shows pools continuously washed through by waves, without Littorinidae, followed by pools only rarely




Figure 3

Beachrock-shore on the leeward side of Curaçao

reached by a wave or only by splash, containing dense populations of *Littorina mespillum* von Mühlfeld, 1824. Members of this species are generally only found below the water line. At the water line of the more landward pools and also near pools with more saline water the 2 species *Nodilittorina tuberculata* and *L. jamaicensis* C. B. Adams, 1850, are common. The ones living at the edge or in the water of the pools usually have extensive erosions of the upper part of their spire. Others, living on rock surfaces or in more dry surroundings have well preserved shells. *Littorina ziczac* populations settle in crevices protected from the sun and the undersides of large boulders that are moistened by splash and spray.

III. Nassau (Figure 4)

A consolidated oolith bar at theseaward side of Paradise Island, Nassau, Bahamas, harbored a number of Littorinidae collected here in the fall of 1970. At the time of collection (1 week) the top of the bar was completely dry. The uppermost parts of the bar were settled by numerous



Figure 4

Transect through a limestone bar on Paradise Island, Nassau

individuals of *Tectarius muricatus* and *Echininus nodulosus* (Pfeiffer, 1832). Near the uppermost spray pools, which were in part filled with brackish water due to rain or were in part dry, *Nodilittorina dilatata* (d'Orbigny, 1841) was common. This population generally occurred in a distinctly lower zone than the former two species. Tidal pools filled with splash water refreshed by daily floods and strong waves contained numerous *L. mespillum*. Rocks from there down into the high water line were settled by *Littorina lineolata* d'Orbigny, 1840.

IV. Fuerteventura (Figure 5)

The Littorinidae of the Canary Islands include 3 species: *Littorina punctata* Gmelin, 1791; *L. neritoides* Linnaeus, 1758; and *L. striata* King & Broderip, 1832. The




Figure 5

Transect of the Islote, a rock projection of the Island of Fuerteventura, Canary Islands

transect shown here was made at the Islote, a large volcanic rock projection extending into the Atlantic Ocean at the west side of the southern part of the island Fuerteventura (Jandia) near the village of Cofete. The powerful waves cause a continuous wetting of rocks above the high tide level and at low tides during most of the year. *Littorina striata* is found from continuously wet spots up into often dry rock areas considerably above the high tide line. It may also be found in spray pools. Here it is often encountered with *L. punctata*. *Littorina neritoides* prefers areas that are high up and only moistened at high tides or if strong winds blow up splash.

On the eastern beaches of the island, more protected from wave action, tidal pools will often contain *Littorina punctata*, while *L. neritoides* prefers crevices and rock surfaces up to the high water line. This species is the marine prosobranch that ventures highest up on the beach in this area. In pebble zones it settles on larger blocks just barely reached by the sea at normal high tides.

V. Banyuls (Figure 6)

Littorina neritoides is the only representative of the Littorinidae commonly found on rock pilings, breakwaters, and cliffs everywhere in the Mediterranean Sea. It was collected from a breakwater sheltering the yacht harbor of the town of Banyuls sur Mer, a small resort town in




Figure 6

Transect across the breakwater of the Banyuls yacht harbor, southern France

southern France near the Spanish border. The periwinkle population is found well above the high tide mark in the spray and splash zone. The largest individuals occur furthest up and are more often seen on flat rocky surfaces, while smaller individuals prefer crevices in rocks. Very heavy seas sometimes sweep over the breakwater. Then *L. neritoides* populations may move upward. Animals trapped in small spray pools on top of the breakwater after the sea has calmed down will die if the water evaporates, leaving the snails surrounded by saline water or dry salt. The animals feed on minute algae that grow on the rocks; they move about and feed only when the surface is moist. Individuals will survive in the absence of moisture for at least 5 months (FRETTER & GRAHAM, 1962). A very detailed study on the ecology of this species was made by LYSAGHT (1941) on the Plymouth breakwater.

VI. Oosterschelde (Figure 7)

At the dikes and on the intertidal flats of the Oosterschelde, a fully marine embayment of the North Sea in the Rhein Delta area of the Netherlands, 3 littorinids can usually be found in great abundance. The dikes are coated with basalt columns on the seaward side up to a level well above the normal high tide line. *Littorina saxatilis* Oliv., 1792 lives just above the high water line in the zone where only splash and spray at normal high tides prevent the growth of grass. This region coincides with the line of




Figure 7

Transect across dike and intertidal area of the Oosterschelde, Netherlands

Explanation of Figures 30 to 35

Figure 30: *Littorina angulifera* from Curaçao × 170
 Figure 31: *Littorina angulifera* from Santa Marta. × 275
 Figure 32: *Littorina nebulosa* from Santa Marta. × 375

Figure 33: *Littorina nebulosa* from Santa Marta. × 375
 Figure 34: *Littorina mespillum* from Curaçao. × 830
 Figure 35: *Littorina mespillum* from Curaçao. × 930



high water spring tides (*cf.* MOORE, 1940 for British occurrence). Consequently, the rock surface is covered by fine algal growths extending down into cavities and crevices between the basalt columns. *Littorina saxatilis* usually is inactive at ebb times, but moves about and feeds when its habitat is moist. The animals are as active in slight rain as when their habitat is moistened by splash water. They feed on algal detritus and diatoms (FRETTER & GRAHAM, 1962).

Usually some individuals of this species will also enter the uppermost *Fucus*-zone, beginning just below the normal high tide line. Here the first individuals of *L. obtusata* Linnaeus, 1758 were found and also the first *L. littorea* Linnaeus, 1758. In the *Fucus* growth, which covers the whole rocky surface of the dike downward to the sand or mud bottom, *L. obtusata* is common, in lower parts with an increasing intermixture of *L. littorea*. *Littorina obtusata* feeds on the tissue of *Fucus* and anything that may have settled there (FRETTER & GRAHAM, 1962).

Littorina obtusata is restricted to rock and to algal growths on hard substrates between mid-tide and mean high water neaps (MOORE, 1940). *Littorina littorea* ranges from the high water mark down to the lowest low tide level, often a bit lower. Its upper limit of distribution can be extended by splash or by artificial sea water creeks as are commonly encountered in Yerseke at the western shore of the Oosterschelde, associated with lobster and oyster culture. *Littorina littorea* lives on rock and among small stones, on gravel or on wooden structures, even on soft sand and mud, but only if stones, boulders, tufts of weed or mussel (*Mytilus edulis* Linnaeus, 1758) clusters provide a firm base in these soft surroundings. Single old specimens may be found regularly below lowest low water, but in less dense populations than above the normal low water line. The limit of rich populations of this species can be observed in the Oosterschelde tidal flats in the presence or absence of *Ulva* growths (presence or absence of green coloration) on mussel beds and boulders. Where *L. littorea* is common, *Ulva* cannot grow, because this alga is eaten by the snails. Washed-in *Ulva* leaves attract the individuals of *L. littorea* during the first hour after the bottom has become dry again, so that up to 1000 animals may be counted in an area of 20 cm × 20 cm, all feeding on *Ulva*. Usually, all *Ulva* parts washed into the area with dense *L. littorea* populations during high tide time will be eaten completely before the arrival of the next high tide. This species also settles on rocks at the water line in brackish water like the Veerse Meer, a brackish water extension of the Oosterschelde without tidal fluctuations.

DESCRIPTION OF THE SHELL

Descriptions of *Littorina littorea*, *L. obtusata* and *L. saxatilis* were given by ABBOTT (1954), BEQUAERT (1943) and ZIEGELMEIER (1966). *Tectarius muricatus* was described by CLENCH & ABBOTT (1942), KAUFMANN & GÖTTING (1970) and well illustrated by ROSEWATER (1972). Descriptions and illustrations of *L. nebulosa* were published by BEQUAERT (*op. cit.*), KAUFMANN & GÖTTING (*op. cit.*), WARMKE & ABBOTT (1962), as well as descriptions of *L. meleagris*. *Littorina angulifera* was described and illustrated by BEQUAERT (*op. cit.*), MARCUS & MARCUS (1963), illustrated by ROSEWATER (*op. cit.*). *Littorina mespillum* was well documented by BEQUAERT (*op. cit.*), ABBOTT (*op. cit.*), WARMKE & ABBOTT (*op. cit.*) *Littorina punctata*, *L. striata*, and *L. neritoides*, with descriptions and drawings, were documented by NORDSIECK (1968). *Echininus nodulosus* was described and illustrated by CLENCH & ABBOTT (*op. cit.*) and in ROSEWATER (*op. cit.*).

The shells of individuals belonging to the *Littorina ziczac* and *Nodilittorina tuberculata* groups are described and illustrated (Figures 8 to 13) again here, to avoid confusion and to define sharply the 3 rediscovered or newly discovered species. This is necessary because they were placed by former authors studying this group into species combinations.

Littorina ziczac Gmelin, 1791 (Figure 8)

Collected along the Colombian coast from Cartagena to Guajira Peninsula and from Curaçao.

Description: Shell elongate conical, whorls 6 to 7, convex. Apex as a rule little or not eroded (submicroscopically), sutures well marked, smooth. Body whorl of adult one half of total height. Shouldered at the periphery, less distinct than in *Littorina jamaicensis*, *L. lineolata* and *L. sp.* First 2 to 3 whorls smooth, sculpture of remainder with very fine wavy lines, 15 to 19 on early whorls, 35 to 40 on the last whorl up to the periphery, and 25 to 35 below the periphery. The engraved spiral lines are difficult to see with the naked eye and shells seem to be glossy and smooth. Sculpture is not weaker below the periphery. Peripheral shoulder smooth and rounded, set off by a shallow rounded ridge. Aperture pear-shaped. Outer lip with a sharp edge, not thickened within, meeting the body whorl at an angle of about 45°, and channeled inside. Inner lip forming a very thin callus over the body whorl.




Figure 8
Shell of *Littorina ziczac*

Columellar area moderately large and wide, slanting inward, very weakly concave. Umbilicus lacking.

Color variable, but generally light grey, fine wavy or zig-zag radial lines of light reddish-brown to dark reddish-brown on bluish or yellowish-white background. Some individuals show yellowish-white background just below the suture and bluish-white above it. Earliest whorls are dark red brown. Mouth within red brown with 2 white spiral bands, one close to the base, the other between the periphery and the suture. Outer lip white, blotched with reddish brown. Columellar area light brown.

Figured by KAUFMANN & GÖTTING (1970: fig. 34).

Littorina sp. (Figure 9)

Collected along the shores from Cartagena to the Guajira Peninsula in Colombia.

Description: Shell elongate conical, whorls generally a little less convex than in *Littorina ziczac*, 7 whorls. Apex



Figure 9
Shell of *Littorina* sp.

as a rule little or not eroded (submicroscopically), microscopically always showing erosion by minute boring organisms (mostly algae). Specimens found living at the edge of tide pools sometimes decollate. Suture well marked and smooth. Body whorl of adult over one half of total height (2 : 3). In fully grown specimens the body whorl is much more convex than earlier whorls and only slightly shouldered at the periphery. Younger (also adult) specimens are more strongly shouldered.

The first 2 to 3 whorls are smooth; sculpture of remaining whorls with moderately deeply engraved spiral lines, on early whorls 12 to 17 and on the body whorl 14 to 20 up to the peripheral edge, and below the edge 10 to 14 often obsolete spiral lines. Peripheral shoulder in old whorls at the end almost smooth, on young ones forming a blunt ridge. Aperture pear-shaped, outer lip with sharp edge, not thickened within, meeting body whorl at a sharp angle forming a narrow groove. Inner lip forming a very slight callus over body whorl. Columellar area long and moderately wide, barely concave, and inclined inward. Umbilicus lacking.

see correction partly opposite

Explanation of Figures 36 to '44

- | | | | |
|---|--------|--|-------|
| Figure 36: <i>Littorina meleagris</i> from Santa Marta. | × 790 | Figure 40: <i>Littorina punctata</i> from Fuerteventura. | × 400 |
| Figure 37: <i>Littorina meleagris</i> from Santa Marta. | × 1280 | Figure 41: <i>Littorina punctata</i> from Fuerteventura. | × 400 |
| Figure 38: <i>Littorina striata</i> from Fuerteventura. | × 290 | Figure 42: <i>Littorina neritoides</i> from Costa Brava. | × 750 |
| Figure 39: <i>Littorina striata</i> from Fuerteventura. | × 370 | Figure 43: <i>Littorina neritoides</i> from Banyuls. | × 400 |
| Figure 44: <i>Littorina ziczac</i> from Curaçao. | × 390 | | |

*Addition to Atlantic Littorinids
by Klaus Barend - Bonn*

Corrections necessary due to error made by the printer:

Explanation of Figures:

Fig. 23-35 (first two plates) correct

Fig. 36: *Littorina lineolata* from Paradise Island Nassau x 430

Fig. 37: *Littorina lineolata* from Paradise Island Nassau x 430

Fig. 38: *Littorina lineolata* from Paradise Island Nassau x 345

Fig. 39: *Naditittorina* ...

Corrections necessary due to error made by the printer:

Explanation of Figures:

- Fig.23-35 (first two plates) correct
Fig.36: *Littorina lineolata* from Paradise Island Nassau x 430
Fig.37: *Littorina lineolata* from Paradise Island Nassau x430
Fig.38: *Littorina lineolata* from Paradise Island Nassau x 345
Fig.39: *Nodilittorina tuberculata* from Santa Marta x 465
Fig.40: *Nodilittorina tuberculata* from Santa Marta x 490
Fig.41: *Nodilittorina tuberculata* from Santa Marta x360
Fig.42: *Echininus nodulosus* from Paradise Island Nassau x 200
Fig.43: *Echininus nodulosus* from Paradise Island Nassau x200
Fig.44: *Echininus nodulosus* from Paradise Island Nassau x 285
Fig.45: *Littorina ziczac* from Santa Marta x440
Fig.46: *Littorina* sp.from Santa Marta x710
Fig.47: *Littorina* sp.from Santa Marta x 400
Fig.48: *Nodilittorina dilatata* from Paradise Island Nassau x420
Fig.49: *Nodilittorina dilatata* from Paradise Island Nassau x 420
Fig.50: *Littorina jamaicensis* from Curacao x390
Fig.51: *Littorina jamaicensis* from Curacao x 410
Fig.52: *Littorina meleagris* from Santa Marta x 790
Fig.53: *Littorina meleagris* from Santa Marta x 1280
Fig.54: *Littorina striata* from Fuerteventura x 290
Fig.55: *Littorina striata* from Fuerteventura x 370
Fig.56: *Littorina punctata* from Fuerteventura x 400
Fig.57: *Littorina punctata* from Fuerteventura x 400
Fig.58: *Littorina neritoides* from Costa Brava x 750
Fig.59: *Littorina neritoides* from Banyuls x 400

Therefore the Figure numbers given along with the descriptions of the radulae page 105-108 must be changed:

- Littorina littorea* to *Littorina nebulosa* are correct.
Littorina meleagris (Fig.52 and 53)
Littorina mespillum (Fig.34 and 35)
Littorina neritoides (Fig.58 and 59)
Littorina punctata (Fig.56 and 57)
Littorina striata (Fig.54 and 55)
Littorina ziczac (Fig.45)
Littorina sp.(Fig.46 and 47)
Nodilittorina dilatata (Fig.48 and 49)
Nodilittorina tuberculata (Fig.39,40, and 41)
Littorina jamaicensis (Fig.50 and 51)
Littorina lineolata (Fig.36,37, and 38)
Echininus nodulosus (Fig.42,43, and 44)



THE VELIGER

The following corrections to Figure Explanations on pages 98 through 106 and Figure referrals on pages 105 to 108 are presented with the apologies of author and editor.

The editor is particularly disturbed about this occurrence as we have been unable to find an explanation for this mixup; it is disturbing since we do not know what additional measures we must take in the future to prevent a repetition of a similar accident.

We present this page without a page number in order that it may be inserted, at the choice of the member or subscriber, where it will be of greatest assistance.

Band in following p. 98

Figure 36: <i>Littorina lineolata</i> from Paradise Island, Nassau	× 430
Figure 37: <i>Littorina lineolata</i> from Paradise Island, Nassau	× 430
Figure 38: <i>Littorina lineolata</i> from Paradise Island, Nassau	× 345
Figure 39: <i>Nodilittorina tuberculata</i> from Santa Marta	× 465
Figure 40: <i>Nodilittorina tuberculata</i> from Santa Marta	× 490
Figure 41: <i>Nodilittorina tuberculata</i> from Santa Marta	× 360
Figure 42: <i>Echininus nodulosus</i> from Paradise Island, Nassau	× 200
Figure 43: <i>Echininus nodulosus</i> from Paradise Island, Nassau	× 200
Figure 44: <i>Echininus nodulosus</i> from Paradise Island, Nassau	× 285
Figure 45: <i>Littorina ziczac</i> from Santa Marta	× 440
Figure 46: <i>Littorina</i> sp. from Santa Marta	× 710
Figure 47: <i>Littorina</i> sp. from Santa Marta	× 400
Figure 48: <i>Nodilittorina dilatata</i> from Paradise Island, Nassau	× 420
Figure 49: <i>Nodilittorina dilatata</i> from Paradise Island, Nassau	× 420
Figure 50: <i>Littorina jamaicensis</i> from Curaçao	× 390
Figure 51: <i>Littorina jamaicensis</i> from Curaçao	× 410
Figure 52: <i>Littorina meleagris</i> from Santa Marta	× 790
Figure 53: <i>Littorina meleagris</i> from Santa Marta	× 1280

Figure 54: <i>Littorina striata</i> from Fuerteventura	× 290
Figure 55: <i>Littorina striata</i> from Fuerteventura	× 370
Figure 56: <i>Littorina punctata</i> from Fuerteventura	× 400
Figure 57: <i>Littorina punctata</i> from Fuerteventura	× 400
Figure 58: <i>Littorina neritoides</i> from Costa Brava	× 750
Figure 59: <i>Littorina neritoides</i> from Banyuls	× 400

The Figure numbers given with the descriptions of the radulae on pages 105 to 108 must be changed as follows:

<i>Littorina meleagris</i> (Figures 52 and 53)
<i>Littorina mespillum</i> (Figures 34 and 35)
<i>Littorina neritoides</i> (Figures 58 and 59)
<i>Littorina punctata</i> (Figures 56 and 57)
<i>Littorina striata</i> (Figures 54 and 55)
<i>Littorina ziczac</i> (Figure 45)
<i>Littorina</i> sp. (Figures 46 and 47)
<i>Nodilittorina dilatata</i> (Figures 48 and 49)
<i>Nodilittorina tuberculata</i> (Figures 39, 40, and 41)
<i>Littorina jamaicensis</i> (Figures 50 and 51)
<i>Littorina lineolata</i> (Figures 36, 37, and 38)
<i>Echininus nodulosus</i> (Figures 42, 43, and 44)

Color variable, but always wavy or zigzag radial chestnut brown stripes and mostly axial lines and stripes of dark brown, grey and black on a white background. Earliest whorls dark brown. Whorl just apical to the suture white or yellowish-white in ground color, above the next suture or the peripheral edge bluish-grey to white. The axial lines may be absent or strongly developed so that the shell appears almost black. Mouth within mahogany brown with 2 white spiral bands, one close to the base, the other between the periphery and the suture. Outer lip white, blotched with reddish brown. Columellar area light brown.

Illustration by KAUFMANN & GÖTTING (1970: fig. 35) as *Littorina lineolata*. This may be *L. ziczac brasiliensis* Vermeij & Porter, 1971. Certainty cannot be evaluated from the short description, and figures were not presented. Ecological data are identical with those of *L. sp.* A *L. ziczac* described in detail by MARCUS & MARCUS (1963) probably is *L. sp.* and agrees with it in the morphology of radula and spawn.

Littorina jamaicensis C. B. Adams, 1850 (Figure 10)

Specimens collected from rocky beaches of Curaçao.

Description: Shell elongate conical with 6 to 7 flattened whorls. The apex in some individuals not eroded (sub-microscopically), in others decollate (found near tide pools). Suture well marked and smooth. Body whorl of adult slightly over one half of total height (4 : 5), just as




Figure 10

Shell of *Littorina jamaicensis*

flattened as earlier whorls. Strongly shouldered at the periphery, almost at a right angle. First 2 to 3 whorls smooth, sculpture of remaining whorls with deeply engraved spiral lines, 6 to 8 on early whorls, and 9 to 11 on the last whorl up to the periphery, and 6 to 7 from the edge to the base. Sculpture in fully grown individuals sometimes weaker below the periphery and here stronger growth lines. Peripheral shoulder forming a narrow blunt ridge. Aperture pear-shaped with an edge where the peripheral shoulder ends. Outer lip with sharp edge, not thickened within, meeting body whorl at a sharp angle, and forming a narrow, slit-like channel with it. Inner lip forming a very slight callus over the body whorl. Columellar area wide, slanting inward, and smooth. Umbilicus lacking.

Color variable, fine wavy or zigzag radial lines of dark red-brown to black, axial lines and stripes black or dark grey on a yellowish background on the apical whorl and bluish-white background on the body whorl. Earliest whorls uniformly pale reddish-brown. Mouth purplish-dark brown within, with 2 white spiral bands, a prominent one close to the base, the other, often weaker, between periphery and suture. Outer lip white, blotched with dark red-brown. Columellar area light to dark brown.

This is *Littorina lineata* of BORKOWSKI & BORKOWSKI, 1969, and probably *L. lineolata* of ABBOTT, 1964.

Littorina lineolata d'Orbigny, 1840 (Figure 11)

All shells were collected at Paradise Island, Nassau, Bahamas.

Description: Shell short-turriculate with 5 to 6 convex whorls. Apex mostly eroded. Suture well marked, smooth. Body whorl of adult over one half of total height (5 : 8), distinctly shouldered at the periphery. First 2 whorls smooth, sculpture of remaining whorls of deeply engraved spiral lines, 5 to 6 on early whorls, 8 to 9 up to the periphery on the last whorl, 6 to 8 weaker ones below the periphery. Peripheral shoulders forming a strong and broad blunt ridge.

Aperture pear-shaped, from peripheral edge to base often almost straight. Outer lip with sharp edge, not thickened within, meeting body whorl at larger angle than in *Littorina jamaicensis* and *L. sp.* (about 90°), channeled inside. Inner lip forming a strong callus over the body whorl. Columellar area moderately long and wide, inclined inward, barely concave. Umbilicus lacking.

Color not as variable as in *Littorina sp.* and *L. jamaicensis*, fine wavy or zigzag chestnut-brown radial lines crossed at the lower part of each whorl by continuous axial grey lines and lines of grey blotches. Background bluish-white. Earliest whorls uniformly pale reddish-brown. Mouth dark brown within with one white spiral band close to the base.




Figure 11

Shell of *Littorina lineolata*

Outer lip with a very narrow white edge, blotched with reddish-brown. Columellar area red-brown.

This is *Littorina lineolata* as described by BORKOWSKI & BORKOWSKI, 1969.

Nodilittorina dilatata (d'Orbigny, 1841) (Figure 12)

Collected at Paradise Island, Nassau, Bahamas.

Description: Shell conical, broadly turriculate, whorls 7 to 8, the first flattened, later more convex. Apex, as a rule, little or not eroded (submicroscopically). Suture wavy, shallow, but distinct between more convex whorls. First 2 to 3 whorls smooth, sculpture of remaining whorls consisting of axial rows of sharp tubercles, 3 rows on early whorls, 7 to 10 beaded rows on the last whorl, between which are found 0 to 5 rough spiral threads. These threads are crossed by rather coarse growth lines. Different individuals have different beaded rows which are most prominent (2 to 4). Aperture subcircular. Outer lip with sharp edge, not thickened within, meeting body whorl at an angle of over 90°. Inner lip forming a very strong callus continuous with the wide columella and thus forming a wide shelf not inclined toward the aperture. This shelf continues to the upper corner of the aperture. Individuals with more than 6 whorls show a conical pseudo-umbilicus.

Color of the first 3 whorls pale reddish-brown, later whorls




Figure 12

Shell of *Nodilittorina dilatata*

light grey to lead grey with whitish tubercles. Edge of the outer lip white with irregular light brown blotches. Columella and aperture dark brown with one clear white stripe below.

The individuals in figure 122 of CLENCH & ABBOTT (1942) closely resemble the one here described.

Nodilittorina tuberculata (Menke, 1828) (Figure 13)

Collected from Santa Marta Rodadero along the rocky shores of the Guajira Peninsula, and on Curaçao.

Description: Shell elongate conical, whorls 7 to 8, flattened. Apex in individuals may be little or not eroded (submicroscopically) or decollate (especially if collected in or near tide pools). Sutures indistinct, wavy. Body whorl of adult slightly over one half of total height (8 : 9). First 3 whorls smooth, sculpture of remaining whorls of axial bands of tubercles, 2 bands on early whorls, and 7 beaded rows on the last whorl. The 2nd and 3rd row of each whorl usually have the largest nodules. Between the beaded rows there are 1 to 4 rough spiral threads, crossed by fine regular growth lines. Aperture oval. Outer lip with sharp edge, not thickened within, meeting body whorl at an angle of about 45° and not channeled inside. Inner lip forming a moderately thick callus over the body whorl. Columella inclined inward, wide with a shallow longitudinal excavation but not forming a shelf as in *Nodilittorina dilatata*. Umbilicus lacking.




Figure 13

Shell of *Nodilittorina tuberculata*

Color brownish-grey to lead-grey with whitish to light orange tubercles. Often greenish colors due to strong algal growths on or within the outer shell surface. The first 3 whorls are light brown, edge of the outer lip white, columella and aperture dark brown with a white stripe below.

This species is illustrated by ROSEWATER (1970), illustrated and described by KAUFMANN & GÖTTING (1970).

SPAWN

Description

Littorina littorea Linnaeus, 1758

The pelagic egg capsules are disc shaped, about 1 mm in diameter, with a swelling in the center in which 1 to 5 eggs (usually 3 to 5) are accommodated. The peripheral parts of the capsule form a flat flange to the central swelling. Floating egg capsules can be found in the waters of the Oosterschelde from April until October. Eggs and veligers of this species are the most common components of the prosobranch plankton in these waters besides veligers of *Hydrobia ulvae* (Pennant). The capsules give rise to free swimming veligers on the 6th day of development (cf. THORSON, 1946 and FRETTER & GRAHAM, 1962).

Tectarius muricatus (Linnaeus, 1758)

Individuals of this species proved to be always viviparous in the area of Santa Marta. Freshly collected females usually shed fully developed veligers if kept overnight in a jar of fresh sea water. These veligers swim actively and carry a shell with a little more than one whorl. In contrast to individuals from Santa Marta, *T. muricatus* from Bermuda (LEBOUR, 1945), from the Barbados (LEWIS, 1960), and from Florida (BORKOWSKI, 1971) produce lens-shaped pelagic capsules with a diameter of about 0.2 to 0.3 mm and with one surface more convex than the other; it is much wider than it is deep with only one egg in each capsule.

Littorina saxatilis Olivi, 1792

This viviparous periwinkle retains its eggs in the brood pouch during development. Hatched individuals crawl through the genital aperture and live in the same rock crevices as the adult animals (FRETTER & GRAHAM, 1962). The egg cocoons are almost as large as those of *Littorina obtusata* (THORSON, 1946). Of females collected at the dikes of the Oosterschelde usually a few contain young throughout the frost-free period of the year. Some groups of this species produce egg masses like those of *L. obtusata*, and fix them to hard substrates (SESHAPPA, 1947).

Littorina obtusata Linnaeus, 1758

Littorina obtusata usually deposits its gelatinous egg mass on damp, unexposed fronds of *Fucus* in the warm months of the summer in the Oosterschelde and elsewhere (FRETTER & GRAHAM, 1962; THORSON, 1946; HERTLING & ANKEL, 1927). The spawn is flat, long, oval, or kidney-shaped, with 40 to 150 $1\frac{1}{2}$ to 2 mm wide round eggs per mass. The mass is composed of 2 to 3 layers, one above the other, embedded in a resistant jelly. After 3 to 4 weeks of development inside the egg capsule, the young hatch in the crawling stage (THORSON, *op. cit.*).

Littorina angulifera Lamarck, 1822

Individuals freshly collected from normal mangrove beaches and of estuarine areas will secrete mucus containing veligers as well as eggs in all stages of development if kept overnight in a jar with fresh sea water. The eggs are all equal in shape and size (0.12 mm) as the inner egg-coverings of pelagic egg-cases in other periwinkles. The female contains eggs and larvae in its brood pouch until extreme flood tides and high waves in storms are calm. Others will move down to the water level

(LEBOUR, 1945; LENDERKING, 1954; MARCUS & MARCUS, 1963) at times and release larvae. But for many individuals seen living on mangrove branches with no vertical extensions to the water surface an active movement to the water is not possible or only with considerable detours. Here, individuals contain eggs and veligers in their brood pouches and will discharge them when sea water wets them. Eggs develop within 3 days into veliger larvae with a transparent, light brown tinted shell of slightly more than one whorl

Littorina nebulosa Lamarck, 1822 (Figure 14A)

Spawn is shed in mucus strings which dissolve on contact with sea water; from it a large number of saucer-shaped capsules of about 0.4mm diameter is set free. These capsules consist of 2 unequally convex cupolas, that are fused by a ring-shaped lamella at the outer rim. The more concave side carries an additional central swelling. Each colorless, transparent pelagic capsule contains one egg surrounded by an inner transparent spherical covering. There is a gelatinous fluid between the outer capsule and the egg-covering, albumen between inner egg-covering and egg. After 4 to 5 days of development the veliger hatches, carrying a shell with little more than one whorl that has a brown nucleus and is otherwise transparent and colorless. Egg cases of *Littorina nebulosa* are very similar with those of *L. flava* King & Broderip, 1832 from Brasil, which, in contrast to *L. nebulosa*, hatch after 2 days of development (MARCUS & MARCUS, 1963).




Figure 14

A: Egg capsule of *Littorina nebulosa*; B: Egg capsule of *Littorina meleagris*. Scale line 0.1mm

Littorina meleagris Potiez & Michaud, 1838 (Figure 14B)

Freshly collected individuals, if kept in a bowl of sea water overnight, produce many cupola-shaped egg capsules measuring 0.2mm in width. A cupola with 3 steps rises on a flat round basal disc. Where cupola and disc are fused, they give rise to a lamella extending outward and vertical to the plane of the basal disc. Each egg-case contains one olive-white egg with its own spherical covering. After 4 days of development a veliger hatches; its transparent shell has little more than one whorl. The egg cases produced by individuals from Florida are identical with those from Santa Marta (BORKOWSKI, 1971).

Littorina mespillum von Mühlfeld, 1824

The spawn of this species is still unknown. Perhaps egg capsules observed by LEWIS (1960) and possibly mistaken as belonging to *Puperita pupa* (Linnaeus, 1767) were produced by this species. In these capsules a rounded cupola with an intermediate ring rises on a flat basal disc. Each egg capsule contains one egg that hatches after two days.

Littorina neritoides Linnaeus, 1758

The pelagic capsule produced by *Littorina neritoides* is lens-shaped, 0.16mm wide, and contains one embryo per capsule; the egg hatches at the veliger stage (LEBOUR, 1935; LINKE, 1935).

Littorina punctata Gmelin, 1791

Littorina striata King & Broderip, 1832

The spawn of these species is unknown.

Littorina ziczac Gmelin, 1791 (Figures 15A, 15B)

Freshly collected animals, if placed overnight in a bowl of sea water, produce pelagic egg capsules throughout the year. Each capsule is about 0.2mm wide and 0.1mm high, beehive-shaped with a flat round disc at the base and a cupola above it. At the edge between both sides a lamellar collar is developed that is inclined outward. The cupola consists of a lower platform with slightly inclined sides at the base and a convex upper part sculptured with

See corrections posted p. 98

Explanation of Figures 45 to 51

- | | | | |
|--|-------|--|-------|
| Figure 45: <i>Littorina</i> sp. from Santa Marta. | × 710 | Figure 48: <i>Nodilittorina dilatata</i> from Paradise Island, Nassau. | × 420 |
| Figure 46: <i>Littorina</i> sp. from Santa Marta. | × 400 | Figure 49: <i>Littorina jamaicensis</i> from Curaçao. | × 390 |
| Figure 47: <i>Nodilittorina dilatata</i> from Paradise Island, Nassau. | × 600 | Figure 50: <i>Littorina jamaicensis</i> from Curaçao. | × 410 |

Figure 51: *Littorina lineolata* from Paradise Island, Nassau. × 430